

# Frontiers in Gamma Ray Spectroscopy

## FIG18



Contribution ID : 54

## Shapes and collectivity of nuclei around $Z = 50$ studied by Coulomb Excitation

### Content :

Numerous experimental and theoretical studies are currently focused on studying the nuclear shell structure of nuclei far from the line of stability. In particular, the evolution of nuclear properties, for example, the reduced transition probabilities across the  $Z = 50$  isotopic chain, has been examined in detail. Radioactive ion beams yield new experimental results close to the doubly magic  $^{100}\text{Sn}$  and  $^{132}\text{Sn}$ , but very accurate data on the stable mid shell nuclei are also of great relevance for our understanding of nuclear structure. The Te nuclei with  $Z = 52$  lies in the transitional region between the spherical nuclei at  $Z = 50$  and deformed Xe and Ba nuclei. For the mid-shell  $^{120,122,124}\text{Te}$  nuclei the partial level show the expected vibrational-like structure with equal energy spacing between the phonon states [1]. In our previous Coulomb excitation experiment [2] at IUAC, New Delhi, where the particle detectors are in the forward direction,  $B(E2; 0^+ \rightarrow 2^+)$  value in  $^{120}\text{Te}$  was re-measured with a much higher precision to allow a comparison with the predictions of the large scale shell model calculations (LSSM). Based on all experimental findings, including the excitation of higher excited states for  $^{120,122,124}\text{Te}$ , on obtained the best agreement with an asymmetric rotor behaviour. The most sensitive probe to characterise a nuclear excitation is via a measurement of quadrupole moments. Therefore, to further investigate the second-order effects in Coulomb excitation (i.e. diagonal matrix elements) and to find an experimental proof of the deformation in  $^{120}\text{Te}$ , the present Coulomb excitation experiment was performed at Heavy Ion Laboratory (HIL), Warsaw, where the particle detectors are in the backward direction. The resulting transitional and diagonal  $E2$  matrix elements, deduced  $B(E2)$  values and static quadrupole moments were extracted using the least square Coulomb Excitation search code - GOSIA [3]. The measured value for  $Q(2^+)$  provides first experimental proof of the prolate shape of the  $^{120}\text{Te}$  nucleus in the  $2^+$  state [4]. These values are also compared with the results obtained using Davydov-Filippov model assuming  $\beta = 0.18$  and  $\gamma \sim 27^\circ$ .

To further study the evolution of nuclear structure properties and in particular, nuclear deformation, across the  $Z=50$  closed shell,  $^{118}\text{Sn}$  nucleus (isotone of  $^{120}\text{Te}$  with  $N=68$ ) was also investigated. This will allow to understand the contribution of single-particle excitation to the collective motion of nuclei. Recent Total Routhian Surface calculations [5] show the co-existence of spherical ground state band with intruder rotational band, making  $^{118}\text{Sn}$  isotope an excellent candidate to study the phenomena of nuclear shape co-existence. With these motivations in mind, a multi-step

Coulomb excitation experiment using  $^{32}\text{S}$  beam was performed at HIL, Warsaw. The preliminary results obtained from this experiment will also be presented during the conference.

**Primary authors** : Dr. SAXENA, Mansi (Heavy Ion Laboratory, University of Warsaw)

**Co-authors** : Dr. NAPIORKOWSKI, P. (Heavy Ion Laboratory, University of Warsaw) ; Dr. SREBRNY, J. (Heavy Ion Laboratory, University of Delhi) ; Prof. WOLLERSHEIM, H. (GSI Helmholtzzentrum für Schwerionenforschung, Germany) ; Dr. KUMAR, Rakesh (Inter University Accelerator Centre, New Delhi) ; Dr. PRÓCHNIAK, L. (Heavy Ion Laboratory, University of Warsaw) ; Dr. STOLARZ, A. (Heavy Ion Laboratory, University of Warsaw) ; Mr. DUTT, Sunil (Department of Physics, Aligarh Muslim University) ; Dr. KOMOROWSKA, M (Heavy Ion Laboratory, University of Warsaw) ; Dr. WRZOSEK-LIPSKA, K. (Heavy Ion Laboratory, University of Warsaw) ; Dr. MATEJSKA-MINDA, M. (Heavy Ion Laboratory, University of Warsaw) ; Dr. PALACZ, M. (Heavy Ion Laboratory, University of Warsaw)

**Presenter** : Dr. SAXENA, Mansi (Heavy Ion Laboratory, University of Warsaw)

**Session classification** : --not yet classified--

**Track classification** : --not yet classified--

**Type** : Poster