


Baryon evolution in relativistic heavy ion collisions

Sandeep Chatterjee
IISER Berhampur

Free Meson Seminar
TIFR, Mumbai

In collaboration with Tribhuban Parida, PhD scholar from IISER Berhampur


The genesis



arXiv: 1610.00646

FIG. 13. (Color online) Beam energy dependence of directed flow slope for protons in 10-40% centrality Au+Au from the STAR experiment, compared with recent available model calculations [19, 20, 95]. All the experimental data are from Ref. [46] except for one energy point, $\sqrt{s_{NN}} = 14.5$ GeV [90], which should be considered a preliminary measurement. The Frankfurt hybrid model [94] as well as a pure hydro calculation with particle freeze-out at constant energy density [94] both lie above the data and are off-scale at all BES energies.

The genesis




Predictions from hydro with Glauber initial conditions need to evolve baryon charge

NOTE: the observable is dv_1/dy at midrapidity, hence probes breaking of Bjorken ansatz of boost invariance


arXiv: 1610.00646

FIG. 14. (Color online) Beam energy dependence of directed flow slope for protons in 10-40% centrality Au+Au from the STAR experiment, compared with recent hybrid [94] and 3FD [95] model calculations. All the experimental data are from Ref. [46] except for one energy point, $\sqrt{s_{NN}} = 14.5$ GeV [90], which should be considered a preliminary measurement.

Since then.....




arXiv: 2003.05852



arXiv: 2207.04927

What is happening with v_2 ?




Hence, we focus on baryon v_1


First, charged particle v_1

Sign convention of v_1


(a) reaction plane




entropy deposition scheme



$$r_1 < r_2 \rightarrow \rho(r_1) > \rho(r_2)$$



entropy deposited scheme



At a generic point (x, y) on the transverse plane, $N_{\text{part}^+}(x, y) \neq N_{\text{part}^-}(x, y)$ where '+' and '-' refer to positive and negative η directions.



This geometric asymmetry has been utilised in Glauber type initial condition models to break boost invariance in the initial condition that can be further evolved by hydro to yield interesting rapidity dependencies in different observables.

entropy deposited scheme

Broadly, 2 schemes have been studied:



Shifted: assume forward-backward (FB) symmetric deposition by a participant source. $N_{\text{part}}^+(x, y) \neq N_{\text{part}}^-(x, y)$ gives rise to a shifted centre of mass rapidity $y_{\text{cm}}(x, y)$ and this causes a shifted fireball [Hirano, Tsuda 2002](#)

Tilted: assume FB asymmetric deposition by a participant source. $N_{\text{part}}^+(x, y) \neq N_{\text{part}}^-(x, y)$ gives rise to a fireball not aligned along the beam axis, tilted fireball. [Bozek, Wyslciel 2010](#)





charged particle v_1 differentiates

Shifted





Tilted



Bozek, Wyskiel 2010
Parida, SC 2022

Now, baryon v_1

Initial condition





$$\epsilon(x, y, \eta_s) = \epsilon_0 [(N_+(x, y) f_+(\eta_s) + N_-(x, y) f_-(\eta_s)) \times (1 - \alpha) + N_{coll}(x, y) \epsilon_{\eta_s}(\eta_s) \alpha]$$

$$f_{+,-}(\eta_s) = \epsilon_{\eta_s}(\eta_s) \epsilon_{F,B}(\eta_s)$$

$$\epsilon_F(\eta_s) = \begin{cases} 0, & \text{if } \eta_s < -\eta_m \\ \frac{\eta_s + \eta_m}{2\eta_m}, & \text{if } -\eta_m \leq \eta_s \leq \eta_m \\ 1, & \text{if } \eta_m < \eta_s \end{cases}$$

$$\epsilon_B(\eta_s) = \epsilon_F(-\eta_s)$$

Initial condition



$$n_B(x, y, \eta_s)$$

$$N_B [W_+^B(x, y) f_+^B(\eta_s) + W_-^B(x, y) f_-^B(\eta_s)]$$

$$W_{\pm}^B(x, y) = (1 - \omega) N_{\pm}(x, y) + \omega N_{coll}(x, y)$$

$$\int \tau_0 dx dy d\eta n_B(x, y, \eta, \tau_0) = N_+ + N_-$$

The transverse profile for baryon, W^B is usually taken $\sim N_{+, -}$. We have allowed for contribution from N_{coll} to account for scenarios that arise in microscopic models like LEXUS where the rapidity loss of colliding energy depositing source depends on the number of binary collisions, thus having more baryon deposited where N_{coll} is large. **NOTE:** Participant source has rapidity asymmetric profile while collision source has rapidity symmetric profile, thus ω decides relative tilt of baryon wrt matter

Evolution


The hydrodynamic evolution of the baryon conserved charge requires the baryon diffusion coefficient κ_B :

$$\kappa_B = \frac{C_B}{T} n_B \left(\frac{1}{3} \coth(\mu_B/T) - \frac{n_b T}{\epsilon + P} \right) \quad \text{Denicol et al 2018}$$


C_B is to be constrained from data

We take, $n_S = 0$, $n_Q = 0.4n_B$, $\frac{\eta T}{\epsilon + P} = 0.08$, $\zeta = 0$, $\epsilon_f = 0.26 \text{ GeV/fm}^3$


Hybrid approach



Role of ω





Role of ω






V_1 of meson (pion) generated by baryon tilt due to pressure anisotropy via EoS

Role of ω







Does not affect rapidity distribution of net proton yield, however the net proton v_1 or splitting in proton - antiproton v_1 is significantly affected



Comparison to data




Comparison to data










Can we constrain C_B (κ_B)?



Sensitive at larger rapidities (baryon rich),
Need measurement at larger rapidities

There is also hints that at lower
 $\sqrt{s_{NN}} \sim 10$ GeV that midrapidity data itself
could constrain

Possible background to EM field effects?

The splitting of directed flow for identified light hadrons (K and p) and strange baryons (Ξ and Ω) in Au+Au collisions at STAR *

[arXiv:2208.01718](https://arxiv.org/abs/2208.01718)

ASHIK IKBAL SHEIKH (FOR THE STAR COLLABORATION)

Department of Physics, Kent State University,
Kent, OH 44242, USA
Email: asheikh2@kent.edu, ashikhep@gmail.com

Received August 4, 2022


The first measurements for rapidity-odd directed flow of Ξ and Ω in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 27$ and 200 GeV are reported. The coalescence sum rule is examined with various combinations of hadrons where all constituent quarks are produced, such as $K^-(\bar{u}s)$, $\bar{p}(\bar{u}\bar{u}\bar{d})$, $\bar{\Lambda}(\bar{u}\bar{d}\bar{s})$, $\phi(s\bar{s})$, $\bar{\Xi}^+(d\bar{s}\bar{s})$, $\Omega^-(sss)$, and $\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})$. For such combinations, a systematic violation of the sum rule is observed with increasing difference in the electric charge and the strangeness content of the combinations. Measurements are compared with the calculations of A Multi-Phase Transport (AMPT) model and Parton-Hadron String Dynamics (PHSD) model with electromagnetic (EM) field. The PHSD model with EM field agrees with the measurements within uncertainties.

Possible background to EM field effects?

arXiv:2208.01718


Index	Quark mass	Charge	Strangeness	Δv_1 combination
1	$\Delta m = 0$	$\Delta q = 0$	$\Delta S = 0$	$[\bar{p}(\bar{u}\bar{u}\bar{d}) + \phi(s\bar{s})] - [K^-(\bar{u}s) + \bar{\Lambda}(\bar{u}\bar{d}\bar{s})]$
2	$\Delta m \approx 0$	$\Delta q = 1$	$\Delta S = 2$	$[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [\frac{1}{3}\Omega^-(sss) + \frac{2}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
3	$\Delta m \approx 0$	$\Delta q = \frac{4}{3}$	$\Delta S = 2$	$[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
4	$\Delta m = 0$	$\Delta q = 2$	$\Delta S = 6$	$[\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})] - [\Omega^-(sss)]$
5	$\Delta m \approx 0$	$\Delta q = \frac{7}{3}$	$\Delta S = 4$	$[\bar{\Xi}^+(\bar{d}\bar{s}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\Omega^-(sss)]$

Possible background to EM field effects?



arXiv:2208.01718

Possible background to EM field effects?



Large contribution from baryon stopping

Summarising..

A new Glauber based model of initial baryon deposition proposed

Qualitative agreement across beam energies with data on yield, v_1

Helps in estimating background driven by baryon stopping across beam energies in signals of other physics like that of the EM field

To be further constrained from baryon - anti baryon splits in other observables like that of polarisation etc

Baryon diffusion may be constrained by proper treatment of the systematics of the parameter space

Independent Evolution of strangeness, electric charge seems important at $\sqrt{s_{NN}} \sim 10$ GeV.

Thank You